

Andrena neffi (Neff's Miner Bee)

No Photo Available

Taxonomy

- **Class:** INSECTA
- **Order:** HYMENOPTERA
- **Family:** ANDRENIDAE
- **Genus:** *Andrena*
- **Scientific Name:** *Andrena neffi* Larkin, 2004
- **Common Name:** Neff's Miner Bee
- **Synonyms:**
- **Taxonomic Name Source:** Integrated Taxonomic Information System (ITIS). 2008. World Bee Checklist Project (version 03-Oct-2008). Integrated Taxonomic Information System: Biological Names. Online. Available: <http://www.itis.gov>.

Agency Status

- **NMDGF:**
- **Federal Status:**
- **BLM Sensitive:**
- **USFS:**
- **IUCN Red List:** [Not Evaluated](#)
- **Nature Serve Global:** [GNR](#)
- **NHNM State:** S1
- **NM Endemic:** YES

Description

Like other *Andrena*, this species is a small, slender, delicate looking bee. It is smaller than most of its congeners; the females are about 8mm in length. The female is mostly black, with some reddish coloration on certain body parts, such as the parts of the clypeus, labrum, antenna, mandibles, and legs. The species is closely related and similar in morphology to *A. ofella*, though it has a broader facial foveae and internal hairs in only about one-third of the corbicula (Larkin 2004).

Habitat and Ecology

Little is known about the habitats and ecology of this species. The collection sites are at about 2,015 m in elevation, in what is likely xeric foothill habitat characterized as a pine-oak woodland. Vegetation types likely include evergreen oaks, alligator junipers and Mexican pines, with an understory that is typically comprised of perennial grasses (Wood Miller *et al.* 2009). The species has been collected only from cinchweed (*Pectis papposa*) and pollen loads from four females was entirely from Helenieae, a tribe in the family Asteraceae (Larkin 2004). It is likely a specialist on Asteraceae, as nearly all related species are specialists, to some degree, on pollen of Asteraceae.

This species likely nests in the ground, like other species in the genus *Andrena* (Michener 2007). In most cases, members of this genus make individual nests, sometimes in large aggregations (Michener 2007). The nests of most communal species are inhabited by several to many females, and typically have only one entrance, with lateral burrows radiating from the main burrow (Michener 2007). The nest cells, which are at the ends of the lateral burrows, are typically lined with a wax like material, to protect larvae from soil moisture and bacteria (Wilson and Carril 2016). Females fill the bottom of each cell with pollen before inserting a single egg. In most cases, species in the genus *Andrena* produce only one generation per year, with mature adults overwintering in nest cells. Typically, the two sexes emerge at a similar time the following year (Michener 2007). This species appears to be active in the late summer; all known records were taken in the last week of August (Larkin 2004).

Geographic Range:

This species is known only from the Mimbres Mountains, in southwestern New Mexico. It may be endemic to these mountains, though survey effort in the region is far from extensive. Four female specimens were collected, in 2000 and 2002, from two localities in the Mimbres Mountains (Larkin 2004).

Conservation Considerations:

There are no conservation measures in place for this species. Research is needed to better understand the distribution, population size and trend, habitats and ecology, threats and habitat trends.

Threats:

The threats to this species are not well understood, though increased wildfire severity and drought due to climate warming are likely threats. The Mimbres Mountains, also known as the Black Range, is at high risk for catastrophic wildfire. For example, in 2013, a lightning strike near Sawyers Peak, 7.5 km as the crow flies from the type locality of this species, ignited a large wildfire. The fire burned almost 140,000 acres of steep and rugged terrain, ranging from mid-elevation pinyon-juniper scrub to high elevation mixed conifer. While some places within the burn scar were unburned, or burned only moderately, 15% of the area experienced stand replacing fire (US Forest Service 2013). Then, in 2022, the second largest fire in New Mexico history burned over 325,000 acres of wilderness in the Black Range (Gleason 2022). In other species, such high severity burns result in habitat degradation due to reduction in canopy and organic ground cover across large areas of canyon slopes, direct mortality from fire, and indirect mortality from post-fire erosion or debris flows (Wallace 2021). The incidence of stand replacing fires has increased in the western U.S., due to in part to climate warming (Turco *et al.* 2023) but primarily due to the legacy of forest management practices, which have changed forest structure, resulting in shrub encroachment in mid elevation pine-oak and pinyon-juniper woodland and desert grasslands and closed canopy conditions in the higher elevation conifer forests. For example, an assessment of the Upper Mimbres watershed found 95% of the landscape was moderately or highly departed from historical conditions (Wood Miller 2009). In the southeastern US, Hanula *et al.* (2015) found low bee diversity in forests where fire suppression resulted in dense tree stands with high understory shrub cover. Open canopies, with little to no mid-story trees and shrubs and a plant diverse understory are optimal for most bee species.

Climate warming may have other unknown impacts on this species, particularly due to increasing drought conditions. The southwestern U.S. saw its driest 22-year period from 2000 to 2021, since at least 800 CE (the time period used in previous climatic reconstructions) (Williams *et al.* 2022). Droughts are projected to become more prolonged, severe, and common in the region under future climate change scenarios (USGCRP 2018). Drought may negatively impact bee species by reducing floral resource availability (Phillips *et al.* 2017). Bees that are oligolectic, like this one, are generally at higher risk of extinction due to reduced host plant availability, especially under climate change scenarios (Roberts *et al.* 2011) and reduced effective population sizes (Packer *et al.* 2005). Additionally, *Andrena* have been reported to have low reproductive output because of the short adult life span, and a low rate of brood cell provisioning (Danforth *et al.* 2019).

While the scope and magnitude of these threats are uncertain, continuing decline in the quality and availability of habitat is inferred. More research is necessary to determine the impacts of these threats on this species.

Population:

The population size and trend are not known for this species. The species is known from only seven females, collected

at two sites across three collection events. This suggests it may be quite rare and range restricted, though survey effort has not been extensive.

References:

- Turco, M., Abatzoglou, J.T., Herrera, S., Zhuang, Y., Jerez, S., Lucas, D.D., AghaKouchak, A. and Cvijanovic, I.. 2023. Anthropogenic climate change impacts exacerbate summer forest fires in California. *PNAS* 120: (e2213815120).
- Packer, L., Zayed, A., Grixti, J.C., Ruz, L., Owen, R.E., Vivallo, F. and Toro, H.. 2005. Conservation genetics of potentially endangered mutualisms: reduced levels of genetic variation in specialist versus generalist bees. *Conservation Biology* 19: (195-202).
- Danforth, B. N., Mincley, R. L., Neff, J. L., and Fawcett, F.. 2019. The solitary bees: biology, evolution, conservation. Princeton University Press.
- Larkin, L.L.. 2004. Four New Fall Species of *Andrena* from the Southwestern United States (Hymenoptera: Andrenidae). *Journal of the Kansas Entomological Society* 77: (254-268).
- Wood Miller, L., Rich, P. and Schumann, M. . 2009. Upper Mimbres Watershed Landscape Assessment. The Nature Conservancy in New Mexico.
- Wallace, J.E.. 2021. Status and Distribution of Terrestrial Snails in Southwestern New Mexico. *Final Report to: New Mexico Department of Game and Fish Share with Wildlife Program*, Tucson, AZ .
- US Forest Service. 2013. Burned Area Emergency Response (BAER) Team Executive Summary. Black Range, Silver City, and Wilderness Ranger Districts Gila National Forest, Silver City, NM : (13).
- Williams, A.P., Cook, B.I. and Smerdon, J.E. . 2022. Rapid intensification of the emerging southwestern North American megadrought in 2020â€“2021. *Nature Climate Change* 12: (232-234).
- [Gleason, M.. 2022. After burning over 325,000 acres of Gila wilderness, the Black Fire is mostly out. Source New Mexico . https://sourcencm.com/2022/07/27/after-burning-over-325000-acres-of-gila-wilderness-the-black-fire-is-mostly-out/](https://sourcencm.com/2022/07/27/after-burning-over-325000-acres-of-gila-wilderness-the-black-fire-is-mostly-out/)
- USGCRP. 2018. Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment, Volume II. *U.S. Global Change Research Program*, Washington, DC, USA : (1515).
- Hanula, J.L., Horn, S. and O'Brien, J.J.. 2015. Have changing forests conditions contributed to pollinator decline in the southeastern United States?. *Forest Ecology and Management* 348: (142-152).
- Phillips, B.B., Shaw, R.F., Holland, M.J., Fry, E.L., Bardgett, R.D., Bullock, J.M. and Osborne, J.L.. 2017. Drought reduces floral resources for pollinators. *Global Change Biology* 24: (3226-3235).
- Roberts, S.P.M., Potts, S.G., Biesmeijer, K., Kuhlmann, M., Kunin, B. and Ohlemüller, R.. 2011. Assessing continental-scale risks for generalist and specialist pollinating bee species under climate change. *BioRisk* 6: (18-Jan).
- Michener, D.C.. 2007. The Bees of the World. The Johns Hopkins University Press, Baltimore : (953).
- [ITIS. 2024. Integrated Taxonomic Information System \(ITIS\). 2024. https://www.itis.gov/](https://www.itis.gov/)
- Wilson, J.S. and Carril, O.M. 2016. The Bees in Your Backyard: A Guide to North America's Bees. Princeton University Press, Princeton, NJ .

More Information

